

Agronomic aspects of biodiversity targeted management of temperate grasslands in Europe – A review

J. Isselstein¹, B. Jeangros² and V. Pavlu³

¹Institute of Agronomy and Plant Breeding, University of Goettingen, Von-Siebold-Str. 8, 37075 Goettingen, Germany

²Agroscope RAC Changins, Swiss Federal Agricultural Research Station, CH-1260 Nyon, Switzerland

³Research Institute of Crop Production, Prague, Grassland Research Station, Rolnicka 6, CZ-46011 Liberec, The Czech Republic

Abstract. Maintaining and enhancing the biodiversity of the agriculturally utilised area has a high priority in environmental policy worldwide. Temperate grasslands in Europe make an important contribution to the biodiversity of agricultural landscapes. The species and community diversity of grasslands is a result of a traditional extensive grassland management interacting with a broad range of site conditions. Until the early decades of the last century, grassland sites were hardly ameliorated and the agronomic potential was generally low, depending on the fertility of the soils. Later on the production from grassland was markedly improved by regular fertilisation, by liming and by artificial drainage of wet sites. Correspondingly, the stocking rates and the cutting frequency increased. Thus, biodiversity strongly decreased, and unimproved species-rich swards only persisted on a low percentage of the total grassland area. The preservation of the remaining species-rich grassland is a primary goal of nature conservation. The continuation of traditional ways of grassland management that would best preserve biodiversity is often not compatible with the requirements of intensive livestock production. Therefore, this grassland is at risk of being abandoned from agricultural use. There is a need to identify and develop improved management measures that better integrate biodiversity and agronomy targets of species-rich grassland farming. In addition, compensation payments for farmers are required to support grass production on species-rich swards. Apart from the unimproved species-rich grassland, there is an increasing area of now de-intensified species-poor grassland which can be managed to increase biodiversity. Grazing at a low stocking rate seems to have the potential to facilitate the restoration of diverse swards and to support reasonable individual performances of the grazing animals.

Key words: grassland management, biodiversity, species-rich grassland, agronomic potential, grazing

INTRODUCTION

Maintaining and enhancing the biodiversity of European grasslands has been given a high priority in national and supranational policy and research initiatives. In the European Union an increasing amount of money is allocated to support the non-production function of agriculture among which biodiversity is a main target (STOA Publication). Grassland plays an important role in the land use of Europe. It is a main part of agricultural production systems and it strongly supports the livelihood of

farmers. In the EU (EU25) more than 30% of the agriculturally utilised area is covered by grassland (EUROSTAT). There is a large variation between countries from less than 5% of grassland in Finland to more than 75% in Ireland. Site conditions, grassland vegetation, management, and production potential are highly variable. European grasslands make an important contribution to the total biodiversity of the rural landscapes (Nösberger & Rodriguez, 1996). A considerable percentage of plants and animal species mainly occur in open grasslands and are rarely found in other vegetation types. In addition, diverse grasslands are highly valued by society due to their potential for recreation and their attraction for tourism (Schübach et al., 2004). Therefore, grasslands have become of primary interest for society, politics and research. About 60% of the farms in Europe that keep cattle are located in disadvantaged regions (Dabbert & Krimly, 2004); this emphasizes the importance of grassland managed for biodiversity and nature conservation as less favoured areas usually have a higher diversity compared to productive areas.

If biodiversity targets are to be integrated into grassland systems, the effect of biodiversity on agronomic performance and profitability of grassland has to be explored. Usually, farmers perceive the biodiversity targeted management of grassland as a limitation to the potential livestock production. The maintenance and enhancement of biodiversity commonly require a reduction of inputs and result in a loss of herbage yield and quality compared to intensively managed grassland. On the other hand, ecologists claim that preserving and restoring grassland diversity may be beneficial to maintain desirable levels of several ecosystem processes, and may therefore have applications in land management and agriculture (Minns et al., 2001). Whether biodiversity could promote the production function of grassland is however largely unexplored and a matter of debate (Isselstein, 2005).

The aim of this paper is to investigate the agronomic consequences of nature conservation and biodiversity targeted grassland management. First, the recent situation of grassland farming in Europe is briefly outlined. The historical development of species-rich grassland and the following loss of diversity as a result of intensification of grassland management are described. Secondly, the management of unimproved swards with a high nature conservation value and of mesic now de-intensified formerly improved swards is investigated.

Grassland farming in Europe today – general options for integrating agronomy and biodiversity targets

The utilisation of grassland in Europe has seen considerable changes during the last two decades. Whereas the overall agriculturally used area in Europe (EU25) has not show a consistent trend of change, the grassland area has decreased slightly in most countries (EUROSTAT). A much stronger decrease occurred in the livestock numbers. Cattle numbers, including dairy and beef, were reduced by 10% from 1992 to 2002 in the European Union (EU25, EUROSTAT, ZMP) with a much stronger decline in the new member states (32% in EU10) compared to the older member states (6% in EU15). A similar situation is found for sheep numbers. The reduction of cattle numbers led to a decrease in beef production of 6% for the whole EU, and of 26% for the new member states. The number of dairy cattle showed a stronger decrease with more than 20% for the whole EU (15% and 38% for EU15 and EU10 respectively). Although the

number of dairy cows decreased markedly, dairy production in the whole EU countries hardly changed during this period. Again, a difference between old and new member states occurred with a slight increase in dairy production in EU15 and a decrease of 15% in EU10. A similar amount of milk had been produced with fewer dairy cows; the milk yield per cow increased by some 30%, from about 4400 kg per cow and year in 1992 to about 5500 in 2001 (ZMP).

These changes in land use and production figures are related to changes in grassland farming. The eastern and central European countries, formerly belonging the COMECON, changed their agricultural land use when they switched from a state to a market economy. Since 1989 livestock numbers have generally decreased by more than 50% up to 70%. The management intensity of improved grassland used by intensive livestock husbandry was reduced, particularly in less favoured regions. Some of the grasslands were even completely abandoned from agricultural use and are now exposed to successional change. In many central and western European countries, such as in Germany, The Netherlands, Denmark and Sweden, the dairy performance was increased by introducing high digestible forages from arable land (e.g. maize) and concentrates into the rations, mainly at the expense of herbage from grassland. Thus, with a fixed milk quota, the percentage of milk being produced from grass is decreasing and grassland is less and less used for dairy production. If technology of production and animal performances continue to progress at current rates, the grassland area needed for dairy farming and intensive grazing will further decrease in the future, unless the demand for dairy and beef products increases significantly (Feehan et al., 2005).

As dairy production is the most profitable way of grassland utilisation, this development will have a strong impact on the maintenance of grassland in the future. Grassland that was no more used for dairy production during the last years has then, to some extent, been de-intensified and further grazed by suckler and beef cattle. However, the market for beef and consequently the potential of grassland based beef cattle farming is limited in Europe. The situation in eastern European countries has demonstrated that some formerly intensively utilised grasslands may be converted into fallow land directly, when dairy and beef production decrease.

With regard to the maintenance and enhancement of grassland biodiversity, the above outlined situation presumably has the following consequences: For grassland that is no longer needed for intensive dairy cow or beef cattle husbandry, intensive management is unlikely to be continued, for economic reasons. Heavy fertilisation and regular sward renovation will be ceased and stocking rates will be low. This management seems to be in accordance with management measures that are generally recommended to enhance biodiversity. However, unimproved species-rich grassland will be increasingly at risk of being abandoned from any agricultural use. Indeed, in the first place, extensive grazing systems will rely on de-intensified grassland which has a history of agricultural improvement and which is therefore superior to unimproved grassland in terms of herbage yield and quality.

For the farmed land as a whole and for grassland in particular, there is an increasing social awareness of the multifunctional character of farming (Jeangros & Thomet, 2004; Lehmann & Hediger, 2004). Grassland provides multiple benefits to farmers and society and among these benefits, biodiversity receives special attention. Agri-environment schemes have been established to accomplish non-market

performances of grassland farming. It is estimated that at least 20% of the agriculturally used area in the EU is managed under an agri-environment scheme (Rounsevell et al., 2005); in Switzerland, ca. 100.000 ha (ca. 13% of the total Swiss grassland area, alpine pastures excluded) of meadows and pastures are managed according to ecological compensation area schemes (Walter et al., 2004). Presumably, the importance of agri-environment schemes for the continuation of farming on de-intensified and on unimproved species-rich grasslands will increase in the future (Rounsevell et al., 2005) and will therefore be essential to maintain and enhance biodiversity.

Historical background of grassland management and grassland biodiversity

The origin of grassland, its vegetation and plant species composition are closely related to human activities and farming practices that have developed since the Neolithic Age. In the temperate regions of Europe, the distribution of natural grassland was limited to sites where natural forests fail to grow, such as sites above the timberline in mountainous regions, very wet or dry sites in the lowlands, or marshlands close to the sea (Ellenberg, 1996). New studies on the history of grasslands suggest that, apart from these sites, grassland could also develop on fertile sites as a result of megaherbivore grazing (Vera, 2000). Grassland dominated open landscapes could have been created in fluvial plains before man controlled the vegetation through deforestation and introduction of crop and animal husbandry. There is evidence that many of the species-rich grassland communities date back to the bronze and iron ages, when regular pasture management and grazing started to establish (Prins, 1998). Since the Middle Ages, when the human population increased, forests on less fertile sites had also been cleared and replaced by grassland and arable land (Korneck et al., 1998). The large diversity of plant species and communities that is found on semi-natural, unimproved grasslands today, has its origin in the Middle Ages and in the early Modern Era (Speier, 1996; Prins, 1998). It is the result of a broad range of site conditions and traditional extensive management systems (Nösberger & Rodriguez, 1996). Soils were not improved until the first half of the 20th century, the agronomic potential was rather low and grass production was dependent on the natural fertility of sites. Traditional grassland utilisation systems were characterised by a generally low level of nutrient return to the swards and, on grazed grassland, by little adaptation of the stocking rates to the actually available herbage. Communal grazing was a common practice. Swards were underutilised in times of herbage surplus and overgrazed when grass growth was limited. In addition, in the traditionally man-made landscape, sheep, cattle and other livestock were important vectors for plant propagules (Poschold et al., 1998).

At the beginning of the 20th century, diverse, unimproved semi-natural grasslands covered a large percentage of the agriculturally utilised area throughout Europe. Starting from this time, grasslands were agriculturally improved and the management intensified. On acid soils with a low fertility, the pH values of the soils and the soil nutrient contents were raised by liming and regular fertilisation. On wet soils, artificial drainage was introduced to remove surplus water. This process was accompanied and followed by an increased use of fertiliser nitrogen and higher stocking rates and defoliation frequency. Thus, grasslands with a high species and community diversity

were replaced by productive pastures and meadows. The result was a dramatic decrease in biodiversity as the swards of improved grasslands were dominated by few productive forage species with a high competitive strength and less productive species were eliminated. Traditionally managed grasslands lost their function as a feed resource for livestock (Willems, 2001) and were reduced to a low percentage of the agriculturally utilised area (Fuller, 1987; Green, 1990; Poschlod, Schumacher, 1998). The remaining semi-natural grasslands are now highly valued by nature conservationists and, in many countries, their maintenance has received priority in nature conservation activities and in agri-environment schemes (Söderstrom et al., 2001; Kumm, 2003; Dipner, & Madel-Kubik, 2004; Jeangros & Thomet, 2004).

Maintaining species-rich grasslands by agricultural management

The maintenance of unimproved semi-natural grasslands requires that grassland management is not intensified (Schellberg et al., 1999), i.e. soils are not ameliorated, no or little fertilisation is applied, and stocking rates are low. However, a regular defoliation of the swards is necessary. It has been frequently suggested that the best way of maintaining semi-natural grasslands would be the maintenance of traditional management measures (Bakker, 1994; Spatz, 1994; Poschlod & Schumacher, 1998; Prins, 1998; Fischer & Wipf, 2002; Muller, 2002). Consequently, management prescriptions laid down in many agri-environment schemes are based on historical ways of agricultural management, such as late cutting, low stocking rates, or reduced fertiliser use. This demonstrates the general problem of integrating semi-natural grasslands into production systems of modern grassland farming. Traditional grassland management systems had been given up during the last century because they no longer met the requirements of the improved livestock production systems in terms of herbage yield and quality and thus were no longer profitable. Therefore, farmers who continue to manage their grasslands in a traditional extensive way act, in many situations, against economic interests unless they receive subsidies to compensate production losses. The amount of payments necessary to maintain a traditional grassland management depends on the gross margin that can be obtained from agricultural products. The more efficiently a semi-natural grassland is integrated into the production process of the grassland farm, the higher is the economic return by the sale of agricultural products. Thus, the need for subsidies to maintain the nature conservation value of these grasslands would be lower and a larger area could be reached with agri-environment schemes within the limitation of public budgets.

Abandonment of semi-natural grasslands from agricultural use is a major risk for grassland diversity. Many investigations have shown that almost independently of the vegetation type, cessation of grassland management leads to successional change and to a loss of plant species diversity. At a local scale, vegetation succession facilitates the invasion of shrubs, the dominance of tall growing species from later successional stages and the competitive exclusion of species typical for managed grasslands (Krahulec et al., 2001; Moog et al., 2002; Tasser & Tappeiner, 2002; Pykala, 2003; Gaisler et al., 2004; Hejzman et al., 2004; Kahmen & Poschlod, 2004). At the regional scale, succession increases the similarity of different grassland communities and thereby reduces the total species diversity (Dullinger et al., 2003).

Major constraints for a continued agricultural management of semi-natural grasslands are the low herbage yield and quality. In a review on the agronomic potential of semi-natural lowland hay-meadows, Tallowin and Jefferson (1999) showed that the herbage growth rate and harvestable yield of species-rich grasslands were at least 50% lower compared to intensively managed meadows. Similar results were obtained from upland and mountainous grasslands (Peeters & Janssens, 1998; Schmid & Jeangros, 1990). The herbage yield of meadows is an important aspect of profitability. As herbage conservation through silage or hay-making is expensive, a minimum herbage yield per cut should be attained; Spatz (1994) suggested 3 tonnes dry matter per ha. This level is frequently missed by marginal grasslands on acid sandy soils or limestone. The herbage quality is even more important for the potential use of herbage from species-rich grasslands. When the primary growth of a species-rich hay-meadow and that of an agriculturally improved grassland are cut at the same time, the digestibility may be the same (Schellberg et al., 1999) or slightly lower on species-rich swards (Tallowin & Jefferson, 1999), obviously depending on the botanical composition. Such comparison is, however, hardly relevant for the farming practice as unimproved swards are less frequently defoliated. Hence, if the late cut primary growth of diverse swards is compared with early cut herbage from improved swards, a large difference between systems is found. This is particularly relevant for many agri-environment schemes where a late first cut is prescribed. Late cut hay from species-rich swards rarely exceeds an organic matter digestibility of 60% (Daccord, 1990; Stoll et al., 2001). Hay or silage of such a low quality cannot be used in intensive livestock systems, i.e. for dairy cows or fattening bulls. It may, however, be fed to dry suckler cows or growing heifers (Spatz, 1994; Stoll et al., 2001).

Grazing is an interesting alternative to cutting in order to maintain species-rich grasslands. Grazing is cheap compared to cutting and forage conservation, it requires less labour input, and the individual performance of the animals is usually higher with grazing compared to barn feeding of conserved late cut herbage. Although some of the traditionally diverse grasslands have developed through long-term hay-cutting and diversity could be best maintained through the continuation of cutting (Spatz, 1994; Fischer & Wipf, 2002; Stammel et al., 2003), there is some evidence that plant species diversity can also be maintained by extensive grazing (Schlapfer et al., 1998; Stammel et al., 2003; Troxler & Chassot, 2004). Plant communities of formerly species-rich grasslands that have been abandoned from agricultural use can be restored by extensive grazing (Hellström et al., 2003; Pykala, 2003). Such grazing creates a mosaic of short and tall grass patches (Bakker et al., 1983; Isselstein et al., 2003). It was shown that in tall grass patches plant species typical for hay-meadows were able to flower and to set seed which is necessary for the persistence of the species in the sward (Correll et al., 2003). In addition, grazing can maintain and increase community diversity at the landscape or regional level (Dullinger et al., 2003). The re-introduction of communal grazing and the establishment of payment schemes that would allow support of the joint utilisation of nature conservation grassland by a group of farmers could be a promising tool for the future management of species-rich grasslands.

Various investigations with grazing sheep and cattle on species-rich, low productive grasslands have shown that the animal performance per unit area is generally low, depending on the fertility of the sites (Jans & Troxler, 1990). However, it also appeared that the individual animal performance is not necessarily lower as

compared to intensive grassland, even if the average herbage quality is lower (Troxler & Chassot, 2004). Obviously, grazers can maintain a good individual performance through selective grazing when the herbage on offer exceeds the demand (Pavlu et al., 2001). This is characteristic for extensive grazing.

Management of de-intensified grassland to enhance biodiversity

Since the early nineties of the last century, an increasing percentage of grassland has been abandoned from intensive agricultural use. Management has either been changed to support less intensive livestock production such as suckler cow or sheep farming or the grasslands have no longer been utilised at all. This is particularly obvious in less favoured areas in central and eastern Europe. Due to a history of agricultural improvement, the biodiversity and therewith the nature conservation value of these grasslands are relatively low. However, many investigations have been performed to study the possibilities to increase biodiversity through extensive management. With regard to the profitability of farming on de-intensified grassland the production costs have to be low. Measures to improve the productivity of the swards like fertilisation, herbicide application, or reseeding etc. are usually not employed and area rents are generally low. Therefore, grazing rather than cutting and forage conservation seems to be the appropriate option for grassland utilisation due to its low costs. In addition, the individual performance of the livestock gains in importance as compared to the performance per unit area. Thus, the management becomes similar to the situation before the agricultural improvement of the sites during the last century. The conditions for a restoration of species diversity are promising in this respect. This was the starting point for various attempts to develop management systems that meet both agronomic and biodiversity targets. A new challenge in the changing agri-environment is therefore to develop management systems that meet both agronomic and biodiversity targets and that are based on the idea that relatively large grassland areas are to be managed with relatively little stock (Rook et al., 2004b).

Various experiments during the last decade have shown that de-intensification of grassland management does not immediately reverse the process of species loss that occurred during the agricultural improvement of the sites. Species number increases if at all slowly (Jeangros & Bertola, 1997; Bakker & Berendse, 1999; Dyckmans et al., 1999; Jeangros & Thomet, 2004; Marriott et al., 2004). Major constraints to the restoration of plant diversity are a high residual fertility of the soils and a continuing high nutrient availability. On mineral soils, phosphorus availability is obviously a key factor (Janssens et al., 1998; Bakker & Berendse, 1999; Koch & Masé, 2001). Herbage production remains on a relatively high level and the dominating forage species prevent the invasion of lost species and the re-establishment of a diverse vegetation. An important prerequisite for the restoration of diverse swards is the decrease in herbage production (Oomes, 1990; Berendse et al., 1992). Unless drastic techniques of nutrient depletion, such as deturfing of the swards, are employed, the reduction of available phosphorus in the soils and the decrease of herbage production is not readily achieved. Therefore, it has been suggested that new techniques should be explored which support the enhancement of species diversity despite a moderate or high nutrient availability (Critchley et al., 2002).

Similarly important is the availability of propagules of plant species that are not present in the current vegetation (Bakker & Berendse, 1999; van Diggelen & Marrs, 2003). Experiments during recent years have shown that the restoration of diverse grasslands was often not successful even if the productivity of the existing swards had decreased. This was due to a lack of appropriate propagules, either from the soil seed bank or from neighbouring vegetation (Bakker et al., 1996; Bekker et al., 1997; Muller et al., 1998; Coulson et al., 2001). In order to overcome this limitation, the addition of propagules had been suggested (Bakker & Berendse, 1999) and investigated in field experiments (e.g. Hopkins et al., 1999; Jones & Hayes, 1999; Kowarsch et al., 2001; Hofmann & Isselstein, 2004). Propagule additions proved to be successful as long as regeneration niches (Grubb, 1977) were sufficiently provided in the sward (Walker et al., 2004). This is primarily dependent on the botanical composition and the sward structure which in turn are controlled by grassland management. Appropriate grazing systems and cutting dates have to be implemented to provide microsite conditions that facilitate germination and establishment of introduced seeds (Smith et al., 2000; Marriott et al., 2004). Recent research on seedling recruitment in lowland grassland suggests that seedling emergence and seedling survival have different requirements on the microsite conditions. Germination and emergence were enhanced in leniently defoliated swards with a low tiller density or where the sward was partly disturbed by mechanical treatment. Obviously, this provided space for germination and emergence. In contrast to emergence of seeds, their survival was improved by frequent defoliation following emergence. This was due to a reduced competition by the existing vegetation (Hofmann & Isselstein, 2004). A great temporal and spatial heterogeneity of the sward structure with patches of short and tall grasses generally occurs with extensive grazing when the herbage on offer exceeds the demand of the grazing livestock (Correll et al., 2003; Kohler et al., 2004). This is a result of selective grazing due to dietary choices of the grazing animals, of treading, and of the heterogeneous return of nutrients to the swards (Rook et al., 2004a). In addition, grazing has a particular advantage over cutting with regard to biodiversity management: the dispersal of plant species propagules is favoured by grazing livestock (Poschlod et al., 1998). To investigate the potential benefits of extensive grazing for the enhancement of biodiversity, an experiment has been set up in five countries throughout Europe. Biodiversity targeted grassland management with low stocking rates is compared with livestock-production orientated management with moderate stocking rates (Rook et al., 2004b). Preliminary results confirmed other findings that the plant species number does not change rapidly in relation to the grazing treatment (Scimone et al., 2004). However, abundance and diversity of butterflies responded more clearly with higher numbers at the low stocking compared to the higher stocking. The reason is obviously the differences in average sward heights (WallisdeVries et al., 2005).

In agronomic terms, grazing seems to have a clear advantage over cutting in many situations. Late cut hay or silage from de-intensified grassland have frequently been demonstrated to have a poor quality and not to be a suitable forage for livestock with increased energy requirements (e.g. Dyckmans et al., 1999; Isselstein et al., 2001). On extensive pasture, the livestock may achieve an individual performance that is close to intensive systems. However, various authors have shown that the performance per unit area of extensive grazing may decrease to 50 to 70% compared to a moderate or intensive grazing (Hofmann et al., 2001; Fothergill et al., 2001; Pavlu et al., 2001;

Barthram et al., 2002; Isselstein et al., 2004). This is an important result for the prospects of integrating extensive grazing into economically viable livestock systems. Assuming the area costs are low, it can be concluded that the individual performance of the livestock is becoming more important and that by this the economic position of extensive grazing compared to intensive grazing is strengthened.

REFERENCES

- Bakker, J.P. 1994. Nature management in Dutch grasslands. In Haggard, R.J., Peel, S. (eds): *Grassland management and nature conservation. BGS Occasional Symposium No 28*, 115–124. Reading, UK.
- Bakker, J.P. & Berendse, F. 1999. Constraints in the restoration of ecological diversity in grassland and heathland communities. *Trends in Ecology and Evolution* **14**, 63–68.
- Bakker, J.P., de Leeuw, W. & van Wieren, S.E. 1983. Micro-patterns in grassland vegetation created and sustained by sheep grazing. *Vegetatio* **55**, 153–161.
- Bakker, J.P., Poschlod, P., Strykstra, R.J., Bekker, R.M. & Thompson, K. 1996. Seed bank and seed dispersal: important topics in restoration ecology. *Acta Botanica Neerlandica* **45**, 461–490.
- Barthram, G.T., Marriott, C.A., Common, T.G. & Bolton, G.R. 2002. The long-term effects on upland sheep production in the UK of a change to extensive management. *Grass and Forage Science* **57**, 124–136.
- Bekker, R.M., Verweij, G.L., Smith, R.E.N., Reine, R., Bakker, J.P. & Schneider, S. 1997. Soil seed banks in European grasslands: does land use affect regeneration perspectives? *Journal of Applied Ecology* **34**, 129–310.
- Berendse, F., Oomes, M.J.M., Altena, H.J. & Elberse, T. 1992. Experiments on the restoration of species-rich meadows in The Netherlands. *Biological Conservation* **62**, 59–65.
- Coulson, S.J., Bullock, J.M., Stevenson, M.J. & Pywell, R.F. 2001. Colonization of grassland by sown species: dispersal versus microsite limitation in response to management. *Journal of Applied Ecology* **38**, 204–216.
- Correll, O., Isselstein, J. & Pavlu, V. 2003. Studying spatial and temporal dynamics of sward structure at low stocking densities: the use of an extended rising-plate-meter method. *Grass and Forage Science* **58**, 450–454.
- Critchley, C.N.R., Chambers, B.J., Fowbert, J.A., Bogha, A., Rose, S.C. & Sanderson, R.A. 2002. Plant species richness, functional type and soil properties of grasslands and allied vegetation in English Environmentally Sensitive Areas. *Grass and Forage Science* **57**, 82–92.
- Dabbert, S. & Krimly, T. 2004. European grassland farms caught between unfriendly markets, perpetually changing agricultural conditions and environmental demands? *Grassland Science in Europe* **9**, 38–48.
- Daccord, R. 1990. Nährwert von Heu aus artenreichen Wiesen. <The nutritive value of hay from species-rich meadows> *Landwirtschaft Schweiz* **3**, 620–624.
- Dipner, M. & Madel-Kubik, E. 2004. New approaches for preservation and promotion of dry grasslands in the conservation politics of Switzerland. *Grassland Science in Europe* **9**, 85–87.
- Dullinger, S., Dirnbock, T., Greimler, J. & Grabherr, G. 2003. A resampling approach for evaluating effects of pasture abandonment on subalpine plant species diversity. *Journal of Vegetation Science* **14**, 243–252.
- Dyckmans, A., Mack, H. & Weissbach, F. 1999. The effect of grassland extensification on yield, forage quality and botanical composition at different grassland locations. *Landbauforschung Völkenrode SH 206*, 125–139.

- Ellenberg, H. 1996. Die Vegetation Mitteleuropas mit den Alpen. <The vegetation of Central Europe including the Alps>. 5th Edition. Ulmer Verlag, Stuttgart, Germany.
- EUROSTAT, Statistical Service of the European Union
http://epp.eurostat.cec.eu.int/portal/page?_pageid=1090,30070682,1090_33076576&_dad=portal&_schema=PORTAL.
- Feehan, J., Gillmor, D.A. & Culleton, N. 2005. Effects of an agri-environment scheme on farmland biodiversity in Ireland. *Agriculture, Ecosystems & Environment* **107**, 275–286
- Fischer, M. & Wipf S. 2002. Effect of low-intensity grazing on the species-rich vegetation of traditionally mown subalpine meadows. *Biological Conservation* **104**, 1–11.
- Fothergill, M., Davies, D.A. & Morgan, C.T. 2001. Extensification of grassland use in Welsh uplands: sheep performance in years 1-6. *Grass and Forage Science* **56**, 105–117.
- Fuller, R.M. 1987. The changing extent and conservation interest of lowland grasslands in England and Wales: a review of grassland surveys 1930-84. *Biological Conservation* **40**, 281–300.
- Gaisler, J., Hejcman, M. & Pavlu, V. 2004. Effect of different mulching and cutting regimes on the vegetation of upland meadow. *Plant Soil Environment* **50**, 324–331.
- Green, B.H. 1990. Agricultural intensification and the loss of habitat, species and amenity in British grasslands: a review of historical change and assessment of future prospects. *Grass and Forage Science* **45**, 365–372.
- Grubb, P.J. 1977. The maintenance of species-richness in plant communities: the importance of the regeneration niche. *Biological Review* **52**, 107–145.
- Hejcman, M., Pavlu, V. & Gaisler J. 2004. Vegetation structure under unmanaged, grazed and cut grassland in the Giant Mountains, Czech Republic. *Grassland Science in Europe* **9**, 213–215.
- Hellström, K., Huhta, A.-P., Rautio, P., Tuomi, J., Oksanen, J. & Laine, K. 2003. Use of sheep grazing in the restoration of semi-natural meadows in northern Finland. *Applied Vegetation Science* **6**, 45–52.
- Hofmann, M. & Isselstein, J. 2004. Seedling recruitment on agriculturally improved mesic grassland: the influence of disturbance and management. *Applied Vegetation Science* **7**, 193–200.
- Hofmann, M., Kowarsch, N., Bonn, S. & Isselstein, J. 2001. Management for biodiversity and consequences for grassland productivity. *Grassland Science in Europe* **6**, 113–116.
- Hopkins, A., Pywell, R.F., Peel, S., Johnson, R.H. & Bowling, P.J. 1999. Enhancement of botanical diversity of permanent grassland and impact on hay production in Environmentally Sensitive Areas in the UK. *Grass and Forage Science* **54**, 163–173.
- Isselstein, J., Benke, M. & Hermanspahn, C. 2001. Bewirtschaftung von Niedermoorgrünland unter Naturschutzaufgaben. <Management of peat grasslands under nature conservation schemes> *Landnutzung und Landentwicklung* **6**, 264–267.
- Isselstein, J., Correll, O., Strodthoff, J., Zhao, G. & Hofmann, M. 2003. Variability of sward structure and plant species composition of pastures at low stocking rates. *Grassland Science in Europe* **8**, 606–609.
- Isselstein, J., Griffith, B.A., Pradel, P. & Venerus, S. 2004. Effects of livestock breed and stocking rate on sustainable grazing systems: 3. agronomic potential. *Grassland Science in Europe* **9**, 620–622.
- Isselstein, J. 2005. Enhancing grassland biodiversity and its consequences for grassland management and utilisation. In: McGilloway D.A., Grassland: a global resource. Wageningen Academic Publishers, 305–320.
- Jans, F. & Troxler, J. 1990. Weidenutzung und Landschaftspflege an Trockenstandorten mit Mutterkühen oder Schafen. I. Tierische Leistung. *Landwirtschaft Schweiz* **3**, 311–314.

- Janssens, F., Peeters, A., Tallowin, J.R.B., Bakker, J.P., Bekker, R.M., Fillat, F. & Oomes, M.J.M. (1998) Relationship between soil chemical factors and grassland diversity. *Plant and Soil* **202**, 69–78.
- Jeanros, B. & Bertola, C. 1997. Changes during six years in botanical composition, species diversity and productivity of a permanent meadow after cessation of fertilizer application and reduction of cutting frequency. *Grassland Science in Europe* **2**, 75–79.
- Jeanros, B. & Thomet, P. 2004. Multi-functionality of grassland systems in Switzerland. *Grassland Science in Europe* **9**, 11–23.
- Jones, A.T. & Hayes, M.J. 1999. Increasing floristic diversity in grassland: the effects of management regime and provenance on species introduction. *Biological Conservation* **87**, 381–390.
- Kahmen, S. & Poschlod, P. 2004. Plant functional trait responses to grassland succession over 25 years. *Journal of Vegetation Science* **15**, 21–32.
- Koch, B. & Masé, G. 2001. Extensivierung von intensiv bewirtschaftetem Grasland. <Extensification of formerly intensively managed grasslands> *Schriftenreihe der FAL* **39**, 61–68.
- Kohler, F., Gillet, F., Gobat, J.-M. & Buttler, A. 2004. Seasonal vegetation changes in mountain pastures due to simulated effects of cattle grazing. *Journal of Vegetation Science* **15**, 143–150.
- Korneck, D., Schnittler, M., Klingenstein, F., Ludwig, G., Takla, M., Bohn, U. & May R. 1998. Warum verarmt unsere Flora? Auswertung der Roten Liste der Farn- und Blütenpflanzen Deutschlands. <Why does the diversity of the flora decrease? An analyses of the Red List of threatened plant species in Germany> *Schriftenreihe für Vegetationskunde* **29**, 299–444.
- Kowarsch, N., Bonn, S., Hofmann, M. & Isselstein, J. 2001. The effect of grassland management and sward disturbance on seedling emergence and establishment of wildflower species. *Grassland Science in Europe* **6**, 186–189.
- Krahulec, F., Skálová, H., Herben, Z., Hadincová, V., Wildová, R. & Pechácková, S. 2001. Vegetation changes following sheep grazing in abandoned mountain meadows. *Applied Vegetation Science* **4**, 97–102.
- Kumm, K.I. 2003. Sustainable management of Swedish seminatural pastures with high species diversity. *Journal for Nature Conservation* **11**, 117–125.
- Lehmann, B. & Hediger, W. 2004. The contribution of grassland to social benefits of agriculture – an economic analysis. *Grassland Science in Europe* **9**, 105–116.
- Marriott, C.A., Fothergill, M., Jeanros, B., Scotton, M. & Louault, F. 2004. Long-term impacts of extensification of grassland management on biodiversity and productivity in upland areas. A review. *Agronomie* **24**, 447–461.
- Minns, A., Finn, J., Hector, A., Caldeira, M., Joshi, J., Palmborg, C., Schmid, B., Scherer-Lorenzen, M., Spehn, E. & Troumbis A. 2001. The functioning of European grassland ecosystems: potential benefits of biodiversity to agriculture. *Outlook on Agriculture* **30**, 179–185.
- Moog, D., Poschlod, P., Kahmen, S. & Schreiber, K.-F. 2002. Comparison of species composition between different grassland management treatments after 25 years. *Applied Vegetation Science* **5**, 99–106.
- Muller, S. 2002. Diversity of management practices required to endure conservation of rare and locally threatened plant species in grasslands: a case study at a regional scale (Lorraine, France). *Biodiversity and Conservation* **11**, 117–1184.
- Muller, S., Dutoit, T., Alard, D. & Grevilliot, F. 1998. Restoration and rehabilitation of species-rich grassland ecosystems in France: a review. *Restoration Ecology* **6**, 94–101.
- Nösberger, J. & Rodriguez, M. 1996. Increasing biodiversity through management. *Grassland Science in Europe* **1**, 949–956.

- Oomes, M.J.M. 1990. Changes in dry matter and nutrient yields during the restoration of species-rich grassland. *Journal of Vegetation Science* **1**, 333–338.
- Pavlu, V., Gaisler, J. & Auf, D. 2001. Intensive and extensive grazing of heifers in the upland Jizerske hory mountains. *Grassland Science in Europe* **6**, 179–182.
- Peeters, A. & Janssens, F. 1998. Species-rich grasslands: diagnostic, restoration and use in intensive livestock production systems. *Grassland Science in Europe* **3**, 375–393.
- Poschlod, P. & Schumacher, W. 1998. Rückgang von Pflanzen des Grünlandes - Gefährdungsursachen und Handlungsbedarf. <Decrease of grassland plant diversity - Causes and necessary actions> *Schriftenreihe für Vegetationskunde* **29**, 83–99.
- Poschlod, P., Kiefer, S., Tränkle, U., Fischer, S. & Bonn, S. 1998. Plant species richness in calcareous grasslands as affected by dispersability in space and time. *Applied Vegetation Science* **1**, 75–91.
- Prins, H.H.T. 1998. Origins and development of grassland communities in Northwestern Europe. In WallisdeVries, M.F., Bakker, J.P. & Van Wieren, S.E. (eds): *Grazing and conservation management*. Kluwer Academic Publishers, Dordrecht, NL, 55–105.
- Pykala, J. 2003. Effects of restoration with cattle grazing on plant species composition and richness of semi-natural grasslands. *Biodiversity and Conservation* **12**, 2211–2226.
- Rook, A.J., Dumont, B., Isselstein, J., Osoro, K., WallisdeVries, M.F., Parente, G. & Mills, J. 2004a. Matching type of livestock to desired biodiversity outcomes in pastures – a review. *Biological Conservation* **119**, 137–150.
- Rook, A.J., Petit, M., Isselstein, J., Osoro, K., WallisdeVries, M.F., Parente, G. & Mills, J. 2004b. Effects of livestock breed and stocking rate on sustainable grazing systems: 1. Project description and synthesis of results. *Grassland Science in Europe* **9**, 572–574.
- Rounsevell, M.D.A., Ewert, F., Reginster, I., Leemans, R. & Carter, T.R. 2005. Future scenarios of European agricultural land use - II. Projecting changes in cropland and grassland. *Agriculture, Ecosystems & Environment* **107t**, 117–135.
- Schellberg, J., Mösel, B.M., Kühbauch, W. & Rademacher, I.F. 1999. Long-term effects of fertilizer on soil nutrient concentration, yield, forage quality and floristic composition of a hay meadow in the Eifel mountains, Germany. *Grass and Forage Science* **54**, 195–207.
- Schlapfer, M., Zoller, H. & Körner, C. 1998. Influences of mowing and grazing on plant species composition in calcareous grassland. *Botanica Helvetica* **108**, 57–67.
- Schmid, W. & Jeangros, B. 1990. Artenreiche Wiesen der Schweiz und ihr Ertrag. <Species-rich meadows of Switzerland and their herbage yield potential> *Landwirtschaft Schweiz* **3**, 610–619.
- Schüpbach, B., Grünig, A. & Walter, T. 2004. Grassland and landscape aesthetics. *Grassland Science in Europe* **9**, 186–188.
- Scimone, M., Smith, R.E.N., Garel, J.P. & Sahin, N. 2004. Effects of livestock breed and stocking rate on sustainable grazing systems: 4. short-term effects on vegetation. *Grassland Science in Europe* **9**, 623–625.
- Smith, R.S., Shiel, R.S., Millward, D. & Corkhill, P. 2000. The interactive effects of management on the productivity and plant community structure of an upland meadow: an 8-year field trial. *Journal of Applied Ecology* **37**, 1029–1043.
- Söderstrom, B., Svensson, B., Vessby, K. & Glimskar, A. 2001. Plants, insects and birds in semi-natural pastures in relation to local habitat and landscape factors. *Biodiversity and Conservation* **10**, 1839–1863.
- Spatz, G. 1994. Freiflächenpflege. <Management of open landscapes> Ulmer Verlag, Stuttgart, Germany.
- Speier, M. 1996. Paläoökologische Aspekte der Entstehung von Grünland in Mitteleuropa. <Paleoecological aspects of the origin of grassland in Central Europe. *Berichte der Reinhold-Tüxen-Gesellschaft* **8**, 199–219.

- Stammel, B., Kiehl, K. & Pfadenhauer, J. 2003. Alternative management on fens: Response of vegetation to grazing and mowing. *Applied Vegetation Science* **6**, 245–254.
- Stoll, W., Arrigo, Y., Chassot, A., Daccord, R., Kessler, J. & Wyss, U. 2001. Bedeutung artenreicher Wiesen als Futter. <Quality of herbage from species-rich hay meadows> *Schriftenreihe der FAL* **39**, 108–114.
- Tallowin, J.R.B. & Jefferson, R.G. 1999. Hay production from lowland semi-natural grasslands: a review of implications for livestock systems. *Grass and Forage Science* **54**, 99–115.
- Tasser, E. & Tappeiner, U. 2002. Impact of land use changes on mountain vegetation. *Applied Vegetation Science* **5**, 173–184.
- Troxler, J. & Chassot, A. 2004. Effects of stocking rate on steer performance and vegetation patterns on mountain pastures. *Grassland Science in Europe* **9**, 587–589.
- van Diggelen, R. & Marrs, R.H. 2003. Restoring plant communities - Introduction. *Applied Vegetation Science* **6**, 106–110.
- Vera, F.W.M. 2000. *Grazing Ecology and Forest History*. Cabi-Publishing, 528 pp.
- Walker, K.J., Stevens, P.A., Stevens, D.P., Mountford, J.O., Manchester, S.J. & Pywell, R.F. 2004. The restoration and re-creation of species-rich lowland grassland on land formerly managed for intensive agriculture in the UK. *Biological Conservation* **119**, 1–18.
- WallisdeVries, M.F., Tallowin, J.R.B., Dulphy, J.P., Sayer, M. & Diana, E. 2005. Effects of livestock breed and stocking rate on sustainable grazing systems: butterfly diversity and abundance. *Grassland Science in Europe* **10**, 227–230.
- Walter, T., Herzog, F., Birrer, S., Dreier, S., Hunziker, M., Jeanneret, Ph., Lüscher, A., Peter, B., Pfiffner, L. & Spiess, M. 2004. Effects of ecological compensation areas on species diversity in the Swiss grassland – an overview. *Grassland Science in Europe* **9**, 171–173.
- Willems, J.H. 2001. Problems, approaches, and results in restoration of Dutch calcareous grassland during the last 30 years. *Restoration Ecology* **9**, 147–154.
- ZMP, Zentrale Markt- und Preisberichtsstelle für Erzeugnisse der Land-, Forst- und Ernährungswirtschaft GmbH Bonn, Agrarmärkte in Zahlen, Issues 1995–2004.