

Seasonal changes in biochemical and microbiological activity of soil against the background of differentiated irrigation in an apple tree orchard after replantation

K. Styła¹, A. Sawicka²

¹ Research Center for Agricultural and Forest Environment, Polish Academy of Science, 60-809 Poznań, Poland; e-mail: styla.katarzyna@gmail.com

² Department of Agricultural Microbiology, University of Life Science, Poznań, 60-656, Poznań, Poland

Abstract. The objective of the studies included seasonal changes in the activity of enzymes, emission CO₂ and microbiological activity of soil against the background of differentiated irrigation in an apple tree (*Malus domestica*) orchard after replantation. The most important activity of enzymes and the number of microorganisms were confirmed in the seasons of spring and autumn, while the lowest activity was observed in summer. The highest dehydrogenase activity was found in the last year (from 0.43 to 2.42 cm³ kg⁻¹) and the lowest the in second year (from 0.19 to 0.58 cm³ kg⁻¹). A high protease activity was recorded in the last year (from 1.25 to 12.08 mg kg⁻¹) and low in the second year (from 1.65 to 3.47 mg kg⁻¹). The highest urease activity was observed in the first year (from 1.17 to 6.42 μmol g⁻¹) and the lowest in the second year (from 0.74 to 2.82 μmol g⁻¹). High intensity of emission CO₂ was noted in summer. The highest CO₂ emission was found in the last year (from 29.32 to 46.86 mg kg⁻¹) and the lowest in the second year (from 15.35 to 27.95 mg kg⁻¹). The highest number of fungi was found in the soil of the combination without irrigation - W₀. A high number of *Azotobacter*, actinomycetes, proteolytic bacteria, phosphate solubilizing bacteria, enzymes activity and CO₂ emission almost always were observed in the soil in the irrigation combination - W₁ or W₂.

Keywords: soil fatigue, replantation, apple tree orchard, irrigation, enzyme and microbiological activity, CO₂ emission

INTRODUCTION

The replant problem refers to the poor growth of replanted fruit trees on old orchard sites and is caused by biotic and abiotic factors. When the problem is caused by biotic factors, it is referred to as replant disease, caused by fungi, bacteria, actinomycetes, nematodes, and their interactions. Abiotic factors that cause replant problem are phytotoxins, nutrient imbalance, low or high pH, soil structure and drainage, and lack or excess of moisture (Utkhede & Smith, 1994).

Many different aspects of the presence and activities of microorganisms in the soil ecosystem can be measured. However, specific microbial parameters indicate different aspects of soil quality. Together with micro- and macrofauna, microorganisms such as bacteria and fungi are key players in the recycling of carbon decomposing the organic matter, followed by the microbes degrading the carbon compounds. The bulk dimension of this ecosystem process can be measured by soil respiration or organic

matter degradation, which at the same time provides an estimate of microbial activity (Kelting et al., 1998, Winding et al., 2005, Liu et al., 2006).

The objective of the present work was to determine the biochemical activity of soil and the number of microorganisms in different terms depending on the developmental phase and level of irrigation in the soil of an apple tree orchard after replantation.

MATERIALS AND METHODS

Studies were conducted from 2003-2005 in an apple tree orchard after replantation at the Agricultural and Fruit-Growing Experimental Farm in Przybroda of the University of Life Science in Poznań, Poland, on a sandy loamy soil (*Albic Luvisol*). The orchard was established in 1975. In 1988, the first replantation was carried out without any preceding soil preparation. In 1994, a repeated replantation was performed, however, after grubbing up the terrain, a shallow ploughing was applied using 2000 kg CaO/ha. Subsequently, apple trees of Šampion cultivar were planted, and as a pollinator Golden Delicious cv. was used on P 60 rootstock in the spacing of 3.5 x 1.5 m (1900 trees/ha). In the experiment, three irrigation levels were used: W_0 - maintenance of soil moisture at the level of atmospheric precipitations, W_1 - maintenance of soil moisture at the level of -0.03MPa of water potential (60% WHC), and W_2 - maintenance of soil moisture at the level of -0.01MPa of water potential (90% WHC). Within each irrigation level, three fertilization combinations were used, i.e. 65 kg N/ha, 65 kg N/ha and 95 kg/ha K_2O and 130 kg N/ha and 190 kg/ha K_2O . Mineral fertilization was applied in the first decade of April, 4-5 weeks before the blooming of orchard trees. Fertilizers were sown in the form of ammonium nitrate and potassium salt (60%). The herbicide program was used during the vegetation season (10 times) from May to September. The following herbicides were applied: Roundup 360 SL (4.0 l ha⁻¹), Chwastox extra (2.0 l ha⁻¹), Basta SL 150 (4.0 l ha⁻¹), Azotop 50 WP (3.0 l ha⁻¹), Agil 100 EC (1.5 l ha⁻¹). Two control samples were introduced: Nowina (apple orchard with Šampion cv. Earlier, agricultural plants were grown on this soil) and Virgin Soil (where no plants were previously grown, the so-called fallow). Soil samples for analyses were taken from belts of herbicide fallow from the depth of 0-20 cm, in three terms depending on the developmental phase of apple trees in the orchard: intensive growth of apple trees (June), fruiting/ripening (August) and fruit harvest connected with leaf fall (October). Bacterial and fungal population sizes were determined using the standard soil dilution plate method. Dilutions were used for the determination of the number of culturable cells as colony forming units (CFU) and expressed on dry matter soil. In the studied soil samples, the following microorganisms were estimated: number of actinomycetes on the Pochon medium (Grabińska-Loniewska, 1999), bacteria from genus *Azotobacter* (Fenglerowa, 1965) and fungi (Martin, 1950). The number of microorganisms were incubated at 24°C during 3, 4, 5 days. The number of proteolytic bacteria was incubated on the Frazier with gelatine medium at 21°C for 48 h (Rodina, 1968). The phosphate solubilizing bacteria was estimated on medium soil with tricalcium phosphate during 10 days incubation at 28°C (Rodina, 1968). The total number of bacteria was developed on 2% agar medium on

soil extract with K_2HPO_4 during 14 days incubation at 28°C (Löchnis, 1920). Dehydrogenase activity in soils was assayed by reduction of triphenyltetrazolium chloride (TTC) to triphenyl formazone (TPF) and expressed as $cm^3 H_2 24 h^{-1} kg^{-1}$ d.m. soil (Thalmann, 1968). Protease activity was assayed by production tyrosine from casein sodium salt and expressed as mg tyrosine $h^{-1} kg^{-1}$ d.m. soil (Ladd & Butler, 1972). Urease activity was measured by released NH_4 and CO_2 from urea and expressed as μmol urea $h^{-1} g^{-1}$ d.m. soil (Hoffmann & Teicher, 1961). The soil emission CO_2 was measured by absorption method and expressed as $mg CO_2 48 h^{-1} kg^{-1}$ f.m. soil (Gołębiowska & Pędziwilk, 1984), Soil pH's were developed from soil-water (1:2 v/v) suspensions by potentiometer method. The total organic carbon (TOC) was analysed on TOC 5050A equipment produced by Shimadzu, Japan. Obtained results were subject to analysis of variance and differences between mean values were estimated by *Tukey's* test. The mean temperatures and total rainfall data during the period of vegetation of the apple orchard are presented in Table 1.

Table 1. Weather characteristics.

| Months | 2003 | | 2004 | | 2005 | |
|----------------------|------------------|--------------------|------------------|--------------------|------------------|--------------------|
| | Temperature (°C) | Precipitation (mm) | Temperature (°C) | Precipitation (mm) | Temperature (°C) | Precipitation (mm) |
| April | 8.9 | 24.5 | 10.0 | 14.9 | 9.6 | 14.2 |
| May | 16.2 | 14.6 | 13.2 | 46.9 | 14.3 | 68.0 |
| June | 20.0 | 24.6 | 16.7 | 63.8 | 17.5 | 11.5 |
| July | 20.8 | 85.7 | 18.7 | 41.7 | 20.4 | 96.6 |
| August | 20.9 | 14.5 | 21.0 | 41.8 | 17.6 | 52.8 |
| September | 15.2 | 19.8 | 14.9 | 33.3 | 16.7 | 52.7 |
| October | 5.6 | 27.9 | 10.4 | 40.6 | 10.5 | 5.6 |
| Mean | 15.3 | 30.2 | 14.9 | 40.4 | 15.2 | 43.0 |
| Sum of precipitation | - | 211.6 | - | 283.0 | - | 301.4 |

Mean temperature and the sum of rainfall in the vegetation period of 2003 (Apr.-Oct.) amounted to 15.3°C and 211.6 mm respectively. In the second year of sample-taking (2004), the mean temperature was 14.9°C and the sum of rainfalls was 283.0 mm. In the year 2005, the respective values were: 15.2°C and 301.4 mm.

RESULTS AND DISCUSSION

The data in Tables 1, 2 clearly show the highest number of microorganisms in autumn and in spring, while in summer, the number was low. The number of fungi in the following years ranged from 33.58-238.85 cfu kg^{-1} in 2003, from 19.48-182.22 cfu kg^{-1} in the second year and from 16.53-260.01 cfu kg^{-1} in the last year. The number of actinomycetes in the following years ranged from 378.57 to 1490.37 cfu kg^{-1} in 2003, from 132.16 to 1696.59 cfu kg^{-1} in the second year and from 386.41 to 1662.72 cfu kg^{-1} in the last year. The number of *Azotobacter* in the following years ranged from 0 to 9 cfu kg^{-1} in 2003, from 0 to 5 cfu kg^{-1} in the second year and from 2 to 50 cfu kg^{-1} in the

last year. The number of proteolytic bacteria in the following years ranged from 15.17 to 266.12 cfu kg⁻¹ in 2003, from 27.25 to 211.72 cfu kg⁻¹ in the second year and from 70.33 to 184.49 cfu kg⁻¹ in the last year. The number of phosphate solubilizing bacteria in the following years ranged from 224.20 to 985.56 cfu kg⁻¹ in 2003, from 297.18 to 1857.55 cfu kg⁻¹ in the second year and from 28.45 to 1406.40 cfu kg⁻¹ in the last year. The total number of bacteria in the following years ranged from 2810.03 to 4539.35 cfu kg⁻¹ in 2003, from 668.45 to 3541.15 cfu kg⁻¹ in the second year and from 350.25 to 4485.90 cfu kg⁻¹ in the last year. Results of the presented work agree with the results obtained by Bis (2002) and Pacholak et al. (2004). Those authors observed an increase of microorganisms in spring and autumn. Malicki (1980) stressed a distinct drop of bacteria number during summer in comparison with spring and autumn. Smoliński et al. (1997) noticed a high number of microorganisms in autumn in comparison with the spring term.

The activities of dehydrogenase, protease and urease were high in autumn or in spring, but low in summer (Table 4). The highest dehydrogenase activity was found in the last year (from 0.43 to 2.42 cm³ kg⁻¹) and the lowest in the second year (from 0.19 to 0.58 cm³ kg⁻¹). The highest protease activity was recorded in the last year (from 1.25 to 12.08 mg kg⁻¹) and the lowest in the second year (from 1.65 to 3.47 mg kg⁻¹). The highest urease activity was found in the first year (from 1.17 to 6.42 μmol g⁻¹) and the lowest in the second year (from 0.74 to 2.82 μmol g⁻¹). Similar results in the activity of enzymes depending on the sampling time were obtained by Bielińska & Domżał (1998). Those authors found that enzymes are comparatively active at the end of spring, while in summer their activity decreases. In autumn, their activity increases again. Bielińska (2001) believes that the decrease of enzymatic activity in summer may be connected with an intensive uptake of nutritive components by the growing trees. On the other hand, the high enzymatic activity of soil in spring can be explained by optimal temperatures with a sufficient level of moisture (Bielińska & Domżał, 1998). These conditions may have a favourable effect on the development of microflora (Bielińska & Domżał, 1998). On the other hand, in autumn, plant residuals which remained after harvest can be the stimulating factor of the increase of enzyme activity. However, results obtained in our present work are different from those shown by Koper & Piotrowska (1999). According to those authors, enzymes were most active in summer (July, August). Bielińska (2001) associated the intensification of enzyme activity in summer with the increase of root mass in that period, which generated the maximal secretion of enzymes to the rhizosphere directly from the roots, as well as with the bacteria and fungi developing in the root zone.

The highest CO₂ emission was observed in the summers of 2003 and 2005 (Table 4). A high CO₂ emission was found in the last year (29.32 to 46.86 mg kg⁻¹) and low in the second year (from 15.35 to 27.95 mg kg⁻¹). Wojnowska et al. (1993) and Dziadowiec & Kaczmarek (1998) also observed a high respiration rate of soil in summer. Mo et al. (2005) noted that the daily soil CO₂ emission was moderate in late spring, increased sharply in summer and decreased in autumn. According to those authors, the soil temperature exerted principle control on the seasonal and annual variation of soil respiration. According to Vanhala (2002) the soil respiration rates decreased during spring and summer, the minimum values occurring at the end of August.

Table 2. Effect of sampling time and irrigation on the number of fungi (F), actinomycetes (A) and *Azotobacter* (Azb) in the soil of apple tree orchard after replantation in the years 2003–2005 (cfu kg⁻¹d.m. soil).

| Sampling time | Combination of irrigation | 2003 | | | 2004 | | | 2005 | | |
|---------------------------------|---------------------------------|-----------------------|-----------------------|-------------------------|-----------------------|-----------------------|-------------------------|-----------------------|-----------------------|-------------------------|
| | | F x10 ⁶ | A x10 ⁶ | Azb x10 ³ | F x10 ⁶ | A x10 ⁶ | Azb x10 ³ | F x10 ⁶ | A x10 ⁶ | Azb x10 ³ |
| Intensive growth of apple trees | W ₀ | 105.48bc* | 455.58a* | 9ab* | 182.22c* | 856.69bc* | 0a* | 102.52bc* | 510.16a* | 12a* |
| | W ₁ | 38.83a | 771.07ab | 1a | 102.73b | 1141.02c | 0a | 35.12a-c | 386.41a | 2a |
| | W ₂ | 33.58a | 517.44a | 1a | 23.74a | 1696.59d | 5ab | 16.53a | 445.30a | 7a |
| Fruiting/ripening | W ₀ | 130.63c | 692.57a | 1a | 72.73.ab | 132.16a | 0a | 55.61a-c | 558.69a | 2a |
| | W ₁ | 57.35ab | 378.49a | 4a | 57.57ab | 365.92a | 0a | 31.83ab | 436.71a | 11a |
| | W ₂ | 41.05a | 1094.57ab | 0a | 130.66bc | 533.97ab | 0a | 31.53ab | 450.63a | 11a |
| Harvesting of fruits | W ₀ | 238.85d | 1096.30ab | 1a | 78.96ab | 471.66ab | 1a | 260.01d | 592.42a | 6a |
| | W ₁ | 79.38abc | 1490.37b | 29b | 58.11ab | 161.34a | 2ab | 113.77c | 1152.45ab | 50b |
| | W ₂ | 55.34ab | 1484.88b | 8ab | 19.48a | 380.33a | 2ab | 48.09a-c | 1662.72b | 4a |
| Means for level of irrigation | W ₀ | 158.32b | 748.15a | 3a | 111.30b | 486.84a | 0a | 139.38b | 553.76a | 6a |
| | W ₁ | 58.52a | 879.97a | 11a | 72.81ab | 556.09a | 1ab | 60.24a | 658.69a | 21b |
| | W ₂ | 43.32a | 1032.30a | 3a | 57.96a | 870.30b | 2b | 92.05a | 852.88a | 7a |
| Means for sampling time | Intensive growth of apple trees | 59.30a | 581.36a | 4ab | 102.90a | 1231.43b | 2a | 51.39a | 447.46a | 7a |
| | Fruiting/ripening | 76.34a | 721.88a | 2a | 86.99ab | 344.01a | 0a | 39.66a | 482.01a | 8a |
| | Harvesting of fruits | 124.52b | 1357.19b | 12b | 52.18b | 337.78a | 2a | 140.62b | 1135.86b | 20b |

* Means marked with the same letters did not differ significantly at the probability of P=0.05
cfu - colony forming units

Table 3. Effect of sampling time and irrigation on the number of proteolytic bacteria (PB), phosphate solubilizing bacteria (PSB) and the total number of bacteria in the soil of apple tree orchard after replantation in the years 2003-2005 (cfu kg⁻¹d.m. soil).

| Sampling time | Combination of irrigation | 2003 | | | 2004 | | | 2005 | | |
|-------------------------------------|---------------------------------|---------------------|----------------------|----------------------|---------------------|----------------------|----------------------|---------------------|----------------------|----------------------|
| | | PB x10 ⁶ | PSB x10 ⁶ | TNB x10 ⁶ | PB x10 ⁶ | PSB x10 ⁶ | TNB x10 ⁶ | PB x10 ⁶ | PSB x10 ⁶ | TNB x10 ⁶ |
| Intensive growth of apple trees | W ₀ | 50.54a* | 224.20b* | 2755.84a* | 49.98a* | 533.85ab* | 2927.42a-c* | 96.41ab* | 28.45b* | 702.38ab* |
| | W ₁ | 146.49b | 985.56a | 2700.82a | 80.64a | 1157.59bc | 3541.15bc | 129.92ab | 147.66ab | 350.25a |
| | W ₂ | 266.12c | 654.36a-c | 2551.19a | 77.45a | 1853.67c | 6783.39e | 184.49a | 186.57ab | 401.31ab |
| Fruiting/ripening | W ₀ | 15.17a | 348.76bc | 2810.03a | 27.25a | 297.18a | 668.45d | 70.33a | 286.26ab | 1233.33a-c |
| | W ₁ | 39.61a | 458.79a-c | 3507.50a | 63.34a | 941.60ab | 2582.76a-d | 130.67ab | 527.67ab | 660.20ab |
| | W ₂ | 62.51a | 693.05a-c | 3948.51a | 211.72b | 1857.55c | 3992.95c | 125.62ab | 618.05a | 1791.10bc |
| Harvesting of fruits | W ₀ | 25.88a | 730.41a-c | 4539.35a | 35.44a | 325.61ab | 1589.60abd | 180.62a | 611.07a | 2372.71c |
| | W ₁ | 40.04a | 982.85a | 4349.31a | 40.58a | 609.40ab | 1107.41ad | 160.66ab | 1292.46c | 4485.90d |
| | W ₂ | 30.06a | 892.66ac | 3643.21a | 61.21a | 547.21ab | 2884.50a-c | 144.12ab | 1406.40c | 4256.83d |
| Means for combination of irrigation | W ₀ | 30.53a | 434.46b | 3368.41a | 37.56a | 385.55a | 1728.50a | 115.79a | 308.59b | 1436.00a |
| | W ₁ | 75.38b | 809.07a | 3519.21a | 61.52a | 902.86b | 2410.44a | 140.42a | 655.93a | 1832.11ab |
| | W ₂ | 119.57c | 746.69a | 3380.97a | 116.79b | 1419.48c | 4553.61b | 151.41a | 737.01a | 2149.75b |
| Means for sampling time | Intensive growth of apple trees | 154.39a | 621.37ab | 2669.28a | 69.36ab | 1181.70a | 4417.32b | 136.94ab | 120.89a | 484.65a |
| | Fruiting/ripening | 39.10b | 500.20a | 3422.01ab | 100.77b | 1032.11a | 2414.72a | 108.87a | 477.33b | 1228.21b |
| | Harvesting of fruits | 31.99b | 868.64b | 4177.29b | 45.74a | 494.07b | 1860.51a | 161.80b | 1103.31c | 3705.15c |

* Means marked with the same letters did not differ significantly at the probability of P=0.05 cfu colony forming units

Table 4. Effect of sampling time and irrigation on the dehydrogenase (ADh), protease, (AP) urease activity (AU) and emission CO₂ (CO₂) in the soil of apple tree orchard after replantation in the years 2003–2005.

| Sampling time | Combination of irrigation | 2003 | | | | | 2004 | | | | | 2005 | | | | | |
|---------------------------------|---------------------------------|---|---------------------------|----------------------------|--|---|---------------------------|----------------------------|--|---|---------------------------|----------------------------|--|---|---------------------------|----------------------------|--|
| | | ADh (cm ³ kg ⁻¹) | AP (mg kg ⁻¹) | AU (μmol g ⁻¹) | CO ₂ (mg kg ⁻¹) | ADh (cm ³ kg ⁻¹) | AP (mg kg ⁻¹) | AU (μmol g ⁻¹) | CO ₂ (mg kg ⁻¹) | ADh (cm ³ kg ⁻¹) | AP (mg kg ⁻¹) | AU (μmol g ⁻¹) | CO ₂ (mg kg ⁻¹) | ADh (cm ³ kg ⁻¹) | AP (mg kg ⁻¹) | AU (μmol g ⁻¹) | CO ₂ (mg kg ⁻¹) |
| Intensive growth of apple trees | W ₀ | 0.63ab* | 3.11a-c* | 2.48a-c* | 23.31a* | 0.34ab* | 2.36a* | 2.82b* | 20.96a* | 0.48a* | 1.51a* | 2.38a* | 41.36ab* | | | | |
| | W ₁ | 0.97a-c | 6.36b-d | 2.73a-c | 26.35a | 0.58b | 3.47a | 1.53a,c | 23.94a | 0.43a | 2.62ab | 1.73a | 36.09ab | | | | |
| | W ₂ | 0.84a-c | 6.72c,d | 3.81b-d | 24.63a | 0.38ab | 3.27a | 1.76a-c | 24.86a | 0.67a | 1.25a | 1.86a | 37.81ab | | | | |
| Fruiting/ripening | W ₀ | 0.49a | 2.61a | 1.81a,b | 22.22a | 0.58b | 2.03a | 1.65a,b | 15.35a | 0.63a | 4.21ab | 1.98a | 35.97ab | | | | |
| | W ₁ | 0.92a-c | 2.12a | 1.17a | 51.21c | 0.19a | 1.65a | 0.74d | 27.95a | 1.22ab | 8.59bc | 2.42a | 34.60a | | | | |
| | W ₂ | 0.60a,b | 3.40a-c | 1.61a | 45.71bc | 0.38ab | 2.29a | 0.76d | 19.82a | 2.02b | 3.86ab | 2.40a | 46.86b | | | | |
| Harvesting of fruits | W ₀ | 1.53b-d | 2.24a | 5.06d,e | 25.89a | 0.38ab | 1.73a | 1.93a,b | 23.60a | 0.55a | 3.95ab | 4.19b | 34.14a | | | | |
| | W ₁ | 1.70c,d | 7.49d | 4.21cd | 22.91a | 0.27a | 2.04a | 2.08a,b | 22.22a | 2.42b | 10.70c | 3.68b | 29.90a | | | | |
| | W ₂ | 2.19d | 5.26a-d | 6.42e | 27.27ab | 0.37ab | 2.03a | 1.17c,d | 27.50a | 1.62a,b | 12.08c | 3.46b | 32.88a | | | | |
| Means for level irrigation | W ₀ | 0.88a | 2.65b | 3.12ab | 23.81b | 0.44a | 2.04a | 1.95b | 19.97a | 0.55b | 3.22a | 2.85a | 37.16ab | | | | |
| | W ₁ | 1.20a | 5.32a | 2.70a | 33.49a | 0.35a | 2.39a | 1.45a | 24.71a | 1.35a | 7.30b | 2.61a | 33.53a | | | | |
| Means for sampling time | W ₂ | 1.21a | 5.13a | 3.95b | 32.54a | 0.38a | 2.53a | 1.23a | 24.06a | 1.44a | 5.73ab | 2.57a | 39.18b | | | | |
| | Intensive growth of apple trees | 0.81a | 5.40a | 3.00b | 24.76a | 0.34a | 3.03b | 1.86a | 23.25a | 0.52b | 1.79a | 1.99a | 38.42a | | | | |
| Fruiting/ripening | Harvesting of fruits | 0.67a | 2.71b | 1.53a | 39.72b | 0.38a | 1.99a | 1.05b | 21.04a | 1.29a | 5.55b | 2.26a | 39.14a | | | | |
| | Harvesting of fruits | 1.81b | 4.99a | 5.23c | 25.36a | 0.44a | 1.93a | 1.73a | 24.44a | 1.53a | 8.91c | 3.77b | 32.31b | | | | |

* Means marked with the same letters did not differ significantly at the probability of P=0.05

The amount and level of water in the soil exert the basic effect on the growth and development of microorganisms living in it. The highest number of fungi was found in the soil of the combination without irrigation - W_0 (55.61 to 260.01 cfu kg⁻¹) and the lowest number of fungi was observed in W_1 and W_2 combination (from 16.53 to 130.33 cfu kg⁻¹). A high number of *Azotobacter*, actinomycetes, proteolytic bacteria, phosphate solubilizing bacteria almost always were observed in the soil in the irrigation combination - W_1 or W_2 (Table 2, 3). The number of actinomycetes ranged from 132.16 to 1096.30 cfu kg⁻¹ in W_0 combination, from 161.34 to 1490.37 cfu kg⁻¹ in W_1 combination and from 380.33 to 1696.59 cfu kg⁻¹ in W_2 combination. The number of *Azotobacter* ranged from 0 to 12 cfu kg⁻¹ in W_0 combination, from 0 to 50 cfu kg⁻¹ in W_1 combination and from 0 to 11 cfu kg⁻¹ in W_2 combination. The number of proteolytic bacteria ranged from 15.17 to 180.62 cfu kg⁻¹ in W_0 combination, from 39.69 to 160.66 cfu kg⁻¹ in W_1 combination and from 30.06 to 266.12 cfu kg⁻¹ in W_2 combination. The number of phosphate solubilizing bacteria ranged from 28.45 to 730.41 cfu kg⁻¹ in W_0 combination, from 147.66 to 1292.46 cfu kg⁻¹ in W_1 combination and from 186.57 to 1857.55 cfu kg⁻¹ in W_2 combination. The total number of bacteria ranged from 668.45 to 4539.35 cfu kg⁻¹ in W_0 combination, from 350.25 to 4485.90 cfu kg⁻¹ in W_1 combination and from 401.31 to 6783.39 cfu kg⁻¹ in W_2 combination.

According to Bielińska (2001), adequate soil moisture is the basic condition for the action of soil enzymes. Water determines the physiological status of microorganisms and plants, and it is also necessary to maintain soil enzymes in catalytically active status. However, the highest activity of the studied enzymes and CO₂ emission was recorded rather in the irrigation combinations, i.e. in W_1 or W_2 (Table 4). Dehydrogenase activity ranged from 0.19 to 2.42 cm³ kg⁻¹ in irrigation combination (W_1 , W_2) and from 0.34 to 1.53 cm³ kg⁻¹ in W_0 combination. Protease activity ranged from 1.25 to 12.08 mg kg⁻¹ in irrigation combination (W_1 , W_2) and from 1.51 to 4.21 mg kg⁻¹ in W_0 combination. Urease activity ranged from 1.65 to 5.06 μmol g⁻¹ in W_0 combination, from 0.74 to 4.21 μmol g⁻¹ in W_1 combination and from 0.76 to 6.42 μmol g⁻¹ in W_2 combination. CO₂ emission ranged from 15.35 to 41.36 mg kg⁻¹ in W_0 combination, from 22.22 to 51.21 mg kg⁻¹ in W_1 combination and from 19.82 to 46.86 mg kg⁻¹ in W_2 combination. Bielińska (2001) obtained only a significant dependence between the moisture and dehydrogenase activity and further pointed to the fact that dehydrogenases showed a higher sensitivity to environmental stress than phosphatases, ureases and proteases being extracellular enzymes creating complexes with soil colloids.

Soil organic matter affects biochemical, chemical, biological and physical soil properties that control soil microbial activity (Dou et al., 2007). A high pH was observed in autumn and in spring, while in summer, pH was low. The highest pH in summer was recorded in the last year. In the following years, pH ranged from 4.79 to 5.90 in 2003, from 3.42 to 6.38 in the second year and from 4.35 to 6.51 in the last year. The highest pH was found in W_2 combination (from 5.83 to 6.35) and the lowest without any irrigation - W_0 (from 4.37 to 4.93). The highest TOC content has been confirmed in summer in the second and last year. A high content of TOC was found in the last year (from 7.03 to 17.32 g kg⁻¹) and low in the second year (from 5.45 to 8.92 g kg⁻¹). However, the highest TOC content was almost always recorded in irrigation combination - W_1 (from 7.03 to 17.32 g kg⁻¹) (Table 5).

Table 5. Effect of sampling time and irrigation on the pH and the content of the total organic carbon (TOC) in the soil of the apple trees orchard after replantation in years 2003-2005.

| Sampling time | Combinat ion of irrigation | 2003 | | 2004 | | 2005 | |
|------------------------------------|---------------------------------------|---------|------------------------------|---------|------------------------------|--------|------------------------------|
| | | pH | TOC (g kg ⁻¹) | pH | TOC (g kg ⁻¹) | pH | TOC (g kg ⁻¹) |
| Intensive growth of apple trees | W ₀ | 4.76a* | 8.31a* | 5.14ab* | 8.71ab* | 4.35c* | 9.12a* |
| | W ₁ | 5.42a-c | 8.96a | 5.36a-c | 8.92ab | 4.94bc | 9.73a |
| | W ₂ | 5.80bc | 8.18a | 6.38c | 8.47ab | 5.85ab | 7.69a |
| Fruiting/ripering | W ₀ | 4.85ab | 8.32a | 3.42a | 8.91ab | 5.61ab | 14.93b |
| | W ₁ | 4.82ab | 7.89a | 4.76a | 9.43b | 6.31a | 17.32c |
| | W ₂ | 5.90c | 7.87a | 5.37a-c | 8.20ab | 6.71a | 14.06b |
| Harvesting of fruits | W ₀ | 4.79a | 10.15a | 4.55a | 6.93ac | 4.85bc | 9.31a |
| | W ₁ | 5.66a-c | 10.20a | 4.50a | 7.50a-c | 6.22a | 7.03a |
| | W ₂ | 5.79bc | 9.82a | 6.02bc | 5.45c | 6.51a | 7.91a |
| Means for level irrigation | W ₀ | 4.80a | 8.92a | 4.37a | 8.18ab | 4.93b | 11.12ab |
| | W ₁ | 5.30b | 9.02a | 4.87b | 8.62b | 5.82a | 11.36b |
| | W ₂ | 5.83c | 8.62a | 5.92c | 7.37a | 6.35a | 9.89a |
| Means for sampling time | Intensive growth of apple trees | 5.33a | 8.48a | 5.63c | 8.70b | 5.05b | 8.85a |
| | Fruiting/ ripering | 5.19a | 8.03a | 4.51a | 8.85b | 6.21a | 15.44b |
| | Harvesting of fruits | 5.41a | 10.06b | 5.02b | 6.62a | 5.86 | 8.08 |

* Means marked with the same letters did not differ significantly at the probability of P=0.05

High microbiological and biochemical activity in the irrigation combination could consist in the higher concentration of TOC in that object. In the irrigation combination were noted high content of TOC and pH or rather low in the combination without irrigation (W₀), similar to the number of microorganisms and enzyme activity in soil. This work confirms, therefore, the increase of microbiological and biochemical activity of soil in those combinations which at the same time have demonstrated favourable chemical properties of the soil. It is well known that active microbial biomass is strongly associated with the pH values of the soil (Meysner et al., 2006). The obtained significant correlation coefficients confirm also the significant dependences between the number of microorganisms, biochemical activity and pH and the content of TOC (Table 6). Chemical properties in the soil may have modified in a significant way the microbiological and biochemical activity in soil in the apple tree orchard after replantation.

Table 6. Correlation coefficients between pH, the content of total organic carbon (TOC), soil temperature and the biological and biochemical activity.

| Parameters | Chemical properties soil | | |
|-----------------------|--------------------------|---------|------------------|
| | pH | TOC | Soil temperature |
| F | -0.18* | -0.04 | -0.21* |
| A | 0.41* | -0.00 | -0.21* |
| Azb | 0.35* | 0.16** | -0.24* |
| PB | 0.21* | -0.02 | 0.09 |
| PSB | 0.35* | -0.00 | -0.08 |
| TNB | 0.31* | -0.10 | -0.04 |
| ADh | 0.42* | 0.15** | -0.32* |
| AP | 0.47* | 0.13*** | -0.15** |
| AU | 0.20* | 0.06 | -0.49* |
| CO₂ | 0.11*** | 0.06 | 0.13*** |

- significant at P <0.001, ** P< 0.01, *** P<0.05.

CONCLUSIONS

Microbiological and enzymatic activity and CO₂ emission of soil were differentiated in the vegetative season and in the particular years of studies. Periods of high activity of microorganisms, higher enzymatic activity of soil in the successive years of studies and in the particular terms may be connected with an increase of pH values and the content of total organic carbon in the soil. Therefore, microbiological and biochemical activities of soil environment are strongly connected with its reaction to and the content of organic matter in soil. Irrigation had a favourable effect on the increased number of typical bacteria, actinomycetes, and biochemical activity while an unfavourable effect of irrigation was exerted on the number of fungi in the soil.

REFERENCES

- Bielińska, E.J. & Domżał, H. 1998. Effect of orchard soil acidification on its biochemical activity. *Zesz. Probl. Post. Nauk Roln.* **456**, 497-502 (in Polish).
- Bielińska, E.J. 2001. Enzymatic activity of soil in sour-cherry orchard depending on cultivation method. *Rozprawy Naukowe Akademii Rolniczej w Lublinie.* **251** (in Polish).
- Bis, H. 2002. Occurrence of toxinogenic fungi in soil environment. In: *Activity of microorganisms in different environments*, Kraków, 35-42 (in Polish).
- Dou, F., Wright, A.L. & Hons F.M. 2007. Depth distribution of soil organic C and N after long-term soybean cropping in Texas. *Soil & Tillage Research*, **94**, 530-536.
- Dziadowiec, H. & Kaczmarek, J. 1998. Effect of stand species composition on the total biological activity of rusty soils in fresh mixed wood". In: *Ecological aspects of soil microbiology*, Poznań, 119-124 (in Polish).
- Fenglerowa, W. 1965. Simple method for counting *Azotobacter* in soil samples. *Acta Microbiol. Polon.* **14**, 203.
- Gołębiowska, J. & Pędziwilk, Z. 1984. CO₂ release as an index of biological activity of cultivated soils. *Acta Microbiol. Polon.* **33**, 249-256.
- Grabińska-Łoniewska, A. 1999. *Laboratory Exercises in General Microbiology*. Oficyna Wydawnicza Politechniki Warszawskiej. Warszawa, **223** (in Polish).
- Hoffmann, G. & Teicher, K. 1961. Ein Kolorimetrisches Verfahren zur Bestimmung der Ureaseaktivität im Boden. *Z. Pflanzenernähr. Dung. Bodenkunde.* **95**, 55-63.
- Kelting, D.L., Burger, J.A. & Gerry, S.E. 1998. Estimating root respiration, microbial respiration in the rhizosphere, and root-free soil respiration in forest soils. *Soil Biol. Biochem.* **39**, 961-968.
- Koper, J. & Piotrowska, A. 1999. Enzymatic activity of soil as a parameter of its fertility evoked by the cultivation system. *Zesz. Probl. Post. Nauk Roln.* **467**, 127-134 (in Polish).
- Ladd, N. & Butler, J.H.A. 1972. Short-term assays of soil proteolytic enzyme activities using proteins and dipeptide derivatives as substrates. *Soil Biol. Biochem.* **4**, 19-30.
- Liu, H.S., Li, L.H., Hang, G.X., Huang, J.H., Sun, J.X. & Wang, H.Y. 2006. Respiratory substrate availability plays crucial role in the response of soil respiration to environmental factors. *Applied Soil Ecology*, **32**, 284-292.
- Löchnis, F. 1920. *Landwirtschaftlich bakteriologisches Praktikum*. Berlin.
- Malicki, J. 1980. Physical properties of soils and their microbiological analysis. *Post. Nauk Roln.* **3**, 45-70 (in Polish).
- Martin, J.P. 1950. Use of acid, rose bengal and streptomycin in the plate method for estimating soil fungi. *Soil Science.* **69**, 215-232.
- Meysner, T., Szajdak, L. & Kuś J. 2006. Impact of the farming systems on the content of biologically active substances and the forms of nitrogen in the soils. *Agronomy Research*, **4** (2), 531-542.
- Mo, W., Lee, M.S., Uchida, M., Inatomi, M., Saigusa, N., Mariko, S. & Koizumi, H. 2005. Seasonal and annual variations in soil respiration in a cool-temperate deciduous broad-leaved forest in Japan. *Agricultural and Forest Meteorology.* **134**, 81-94.
- Pacholak, E., Zydlik, Z. & Sawicka, A. 2004. Effect of fertilization on the microbiological status of soil in a replanted apple-tree orchard. Part II. Number of bacteria. *Prace Kom. Nauk Rol. i Kom. Nauk Leś. PTPN.* **97**, 307-316 (in Polish).
- Rodina, A. 1968. *Microbiological studies of waters*. Wydawnictwo Rolnicze i Leśne. Warszawa (in Polish).
- Smoliński, S., Kotwica, K., Jaskulski, D. & Tomalak, S. 1997. Effect of stubble crop on microbiological activity of soil. Changes in the number of bacteria participating in the transformations of C and N. In: *Microorganisms in environment. Occurrence, activity and importance*, Kraków, 625-630 (in Polish).

- Thalman, A. 1968. Zur Methodik der Bestimmung der Dehydrogenase Aktivität in Boden Mittels Triphenyltetrazoliumchlorid (TTC). *Landwirtsch. Forsch.* **21**, 249-258.
- Utkhede, R.S. & Smith, E.M. 1994. Biotic and abiotic causes of replant problems of fruit trees. *Acta Horticulturae*, **363**, 25-31.
- Winding, A., Hund-Rinke, K. & Rutgers, M. 2005. The use of microorganisms in ecological soil classification assessment concepts. *Ecotoxicology and Environmental Safety*. **62**, 230-248.
- Wojnowska, T., Sienkiewicz, S. & Wojtas, A. 1993. Dynamics of CO₂ emission from soil depending on fertilization with manure and NPK as well as on plant cultivation. *Zesz. Probl. Post. Nauk Roln.* **411**, 101-105 (in Polish).
- Vanhala, P. 2002. Seasonal variation in the soil respiration rate in coniferous forest soils. *Soil Biol. Biochem.* **4**, 1375-1379.